

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

A hole in the sky – The dependence of the galaxy luminosity function on the environment

S. Phleps, C. Wolf, J.A. Peacock, K. Meisenheimer, and E. van Kampen

Abstract. We have developed a method to calculate overdensities in multicolour surveys, and compare the local density contrast measured in galaxy samples with different redshift error distributions. We calculated overdensities for three COMBO-17 fields, and identified a region in the CDFS, where the density is lower by a factor of 2 compared to the other two fields. This is mainly due to a deficiency of faint red galaxies. This result is in agreement with local observations in the 2dF.

It is a well established fact that galaxy properties such as morphology and colour depend on environment – red, early-type galaxies are usually found in denser regions of the universe, whereas the field is dominated by blue, late-type galaxies. The dependence of the galaxies' appearance on the local density is a clue to their formation and evolution, so it is important to investigate it in some detail. A powerful means to do that is the type-dependent luminosity function (LF in the following), calculated in different overdensity regimes. With spectroscopic data it is possible to measure the local galaxy density contrast $\delta_g = (\rho/\bar{\rho}_0) - 1$ in small spheres of e.g. $8h^{-1}$ Mpc. However, for an analysis at higher redshifts, deep multicolor surveys have to be used, and those have too large redshift inaccuracies to facilitate a determination of δ_g in small spheres. Instead, δ_g can be calculated in small redshift bins (with binsize $\Delta z = 0.02$ and in steps of $\delta_z = 0.005$). We tested the influence of redshift errors on the determination of δ_g using a COMBO-17 mock survey (van Kampen et al. 2005), and found that the structures are smoothed according to the size of the errors. Red galaxies, which are statistically brighter, have smaller errors than the (statistically) fainter blue galaxies, so in order to directly compare the two samples, we have to calculate a blurring function which renders their redshifts as inaccurate as the blue ones.




Figure 1. **Left:** Overdensities in three COMBO-17 fields. **Right:** For red and blue galaxies in the CDFS. The vertical lines indicate the range in which the LF is calculated.

The blurring function can be found via the convolution theorem and then applied to the data in a Monte-Carlo fashion. We used COMBO-17 data (see Wolf et al. 2003) and calculated the overdensities for all galaxies with $R \leq 23.65$ and rest-frame B -band magnitudes $M_B \leq -18.0$, see left panel of Fig. 1. In the redshift range $0.25 \leq z \leq 0.4$ the CDFS (Chandra Deep Field South) is underdense with respect to the other two fields, where the underdensity fluctuates about the mean. We split the sample into red sequence and blue cloud galaxies (see Bell et al. 2004) and repeated the calculation after blurring the red galaxies' redshifts. In the calculation of the mean density $\bar{\rho}$ the increasing number density of blue galaxies with redshift has to be taken into account, which we estimate from a fit to the measured mean in the three fields. The right panel of Fig. 1 shows the resulting underdensities in the CDFS. The underdensity at $0.25 \leq z \leq 0.4$ is clearly deeper in the red sample. We calculated the type-dependent LF in this redshift range, for each field separately, see Fig. 2. While the LF of




Figure 2. The luminosity function of the red sequence (upper panel) and blue cloud (lower panel) galaxies in the redshift range $0.25 \leq z \leq 0.4$, for the three COMBO-17 fields. The STY fit is overplotted in each panel.

the blue galaxies remains unaffected by environment, the LF of the red galaxies does depend on the local density: In the CDFS the faint-end slope is clearly more positive than in the other fields. Hence, the 'hole in the sky' is mainly due to a deficiency of faint red galaxies. This result is in general agreement with the result of Croton et al. (2005), who found the LF of blue 2dF galaxies to be essentially constant with varying overdensity, and the faint-end slope of the LF of the red galaxies to be steeper in higher density regions.

References

- Bell, E. F., Wolf, C., Meisenheimer, K., et al. 2004, *ApJ*, 608, 752
 Croton, D. J., Farrar, G. R., Norberg, P., et al. 2005, *MNRAS*, 356, 1155
 van Kampen, E., Percival, W.J., Crawford, M. et al. 2005, *MNRAS*, 359, 469
 Wolf, C., Meisenheimer, K., Rix, H.-W., et al. 2003, *A&A*, 401, 73